



Maury, J., Zawodny, W., & Clayden, J. (2017). Stereospecific Intramolecular Arylation of 2- and 3-Pyridyl Substituted Alkylamines via Configurationally Stable α -Pyridyl Organolithiums. *Organic Letters*, 19(3), 472-475. <https://doi.org/10.1021/acs.orglett.6b03603>

Peer reviewed version

Link to published version (if available):
[10.1021/acs.orglett.6b03603](https://doi.org/10.1021/acs.orglett.6b03603)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via ACS at <http://pubs.acs.org/doi/abs/10.1021/acs.orglett.6b03603>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>


Stereospecific Intramolecular Arylation of 2- and 3-Pyridyl Substituted Alkylamines via Configurationally Stable α -Pyridyl Organolithiums

Julien Maury^{‡,a,b}, Wojciech Zawodny^{‡,a} and Jonathan Clayden^{b*}

^a School of Chemistry, University of Manchester, Oxford Road, Manchester M13 9PL, U.K.

^b School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, U.K.


Supporting Information Placeholder



ABSTRACT: Treatment of *N'*-aryl urea derivatives of enantiomerically-enriched α -(2-pyridyl) and α -(3-pyridyl)alkylamines with base leads to the migration of the *N'*-aryl substituent from *N* to *C* in a 'non-classical' intramolecular nucleophilic aromatic substitution reaction. Both electron-rich and electron-poor rings migrate successfully. A new quaternary stereogenic centre is formed adjacent to the pyridine ring with high stereospecificity, even when the intermediate anion is a presumably planar 2-picolylolithium. Base hydrolysis of the urea gives enantiomerically enriched α -pyridylalkylamines.

Pyridines play a vital role in medicinal chemistry,¹ being the most common heterocyclic ring encountered in small molecule drugs.² Substituted chiral pyridines with a stereogenic centre at the 'picolinic' position α to the pyridine ring are present in many biologically active molecules³ and chiral ligands.⁴ More specifically, congested quaternary stereogenic centres bearing both pyridine and phenyl rings are present in antihistamines such as pheniramine⁵ and doxylamine⁶ and in potent cholesteryl ester transfer protein (CETP) inhibitors (Figure 1.).⁷


Figure 1. Bioactive compounds with a quaternary stereogenic centre α to a pyridine ring



Methods for the synthesis of α -chiral amines bearing a pyridine ring at the stereogenic centre typically rely on auxiliaries to direct addition to, or reduction of, an intermediate imine.⁸ For enantiopure α -tertiary amines with a pyridine ring as one of the substituents at the quaternary

stereogenic centre (Figure 1) synthetic approaches are very limited.⁹ We previously reported a stereospecific route to a subclass of these structures by intramolecular migration of pyridine rings to the α -position of lithiated urea derivatives of *N*- α -methylbenzylamine, giving α -tertiary amines after the solvolysis of the urea (Scheme 1).^{9c} Stereospecificity is ensured by the configurational stability of the benzyllithium intermediates¹⁰ on the time scale of the rearrangement reaction.¹¹

Scheme 1. α -Pyridylation of chiral amines.²




We now report a complementary method for the generation of pyridine-bearing quaternary stereogenic centers by stereospecific intramolecular arylation¹² α to 2- or 3-pyridyl substituents. The reaction is mediated by pyridine-stabilised organolithiums that exhibit remarkable

configurational stability, given the electron-withdrawing, anion-stabilising nature of the pyridyl substituents.


Chiral amine precursors **5** and **6** bearing an amino group and a 3- or 2-pyridyl substituent at the stereogenic centre were made using Ellman's *N*-sulfinyl auxiliary.⁸ Grignard addition¹³ and indium allylation¹⁴ of *N*-sulfinyl aldimines **2** or diastereoselective reduction of *N*-sulfinyl ketimines¹⁵ **1** gave highly enantioenriched sulfinamides **3** and **4** that were hydrolysed to chiral primary amines **5** and **6** (Scheme 2). The amines were either acylated (with *N*-methylcarbamoyl chlorides) and methylated, or reductively aminated with *p*-methoxybenzaldehyde and acylated (with aryl isocyanates, followed by methylation, or with *N*-methylcarbamoyl chlorides), giving ureas **7-10** as starting materials for organolithium-mediated rearrangements.

Scheme 2. Synthesis of enantioenriched 2- and 3-pyridyl-substituted urea starting materials. ^aEnantiomeric sulfinimine (*R*)-**2** was used for the synthesis of (*R*)-**5**.



The 3-pyridyl-substituted urea **7a** (R = Me, Ar = Ph) was treated with LDA in THF at -78 °C (Scheme 3). After 3 hours, the reaction was quenched, and rearranged urea **11a**, in which the phenyl group had migrated from nitro to the position α to the pyridyl ring, was isolated in good yield without loss of enantiomeric purity (Table 1, entry 1). No additives¹⁶ were required to maintain the stereospecificity¹⁷ of the reaction, indicating that the presumed intermediate 3-pyridine-stabilised organolithium **7Li** does not racemise on the time scale of the rearrangement. The electron-deficient *para*-fluorophenyl, *para*-chlorophenyl, and 2-pyridyl rings of **7b-d** likewise migrated to give **11b-d** in good yields, again with full stereospecificity (entries 2-4). Urea **11d**, formed in 98:2 er, provides the first example of an α -tertiary amine derivative with both a 2- and a 3-pyridyl substituent at the stereogenic center. The tolerance of the reaction to steric

hindrance¹⁸ was explored by replacing the α -methyl substituent with a benzyl group to build structural analogues of the CETP inhibitor in Figure 1 (entries 5-8). The rearrangements of **7e-7h** were fully stereospecific, and **11e** and **11h** were formed in good yields.



Scheme 3. Stereospecific intramolecular arylation of 3-pyridine-substituted ureas. For simplicity, reactions of the *S* enantiomers are shown; in the case of **7e-h** the *R* enantiomer was used (see Table 1).

Table 1. Arylation of 3-pyridine-substituted ureas


entry	SM, er ^a	R	Ar	product, yield (%)	product, er ^a
1	(<i>S</i>)- 7a 98:2	Me	C ₆ H ₅	(<i>R</i>)- 11a 65	98:2
2	(<i>S</i>)- 7b 98:2	Me	4-FC ₆ H ₄	(<i>R</i>)- 11b 74	98:2
3	(<i>S</i>)- 7c 98:2	Me	4-ClC ₆ H ₄	(<i>R</i>)- 11c 63	98:2
4	(<i>S</i>)- 7d 98:2	Me	2-pyridine	(<i>S</i>)- 11d 76	98:2
5	(<i>R</i>)- 7e >99:1	Bn	C ₆ H ₅	(<i>S</i>)- 11e 87	>99:1
6	(<i>R</i>)- 7f >99:1	Bn	3-ClC ₆ H ₄	(<i>S</i>)- 11f 22	>99:1
7	(<i>R</i>)- 7g >99:1	Bn	3-MeOC ₆ H ₄	(<i>S</i>)- 11g 26	>99:1
8	(<i>R</i>)- 7h >99:1	Bn	2-pyridine	(<i>R</i>)- 11h 88	>99:1

^a Enantiomeric ratio by HPLC on chiral stationary phase.

Hydrolysis of the rearranged products under basic conditions cleaved the urea in good yield to provide α -tertiary amines **12**, illustrated by the formation of **12b** and **12d** (Scheme 3).


The absolute configuration of (*R*)-**11c** was determined by means of X-ray crystallography (Figure 2), and confirmed that the rearrangements of 3-pyridyl ureas proceed with retention of configuration, as has been observed in previously related rearrangements¹¹.

Figure 2. X-ray crystal structure of (*R*)-**11c**.




Replacing the *N*-methyl substituent with an *N*-*p*-methoxybenzyl (PMB) protecting group had no effect on the stereospecificity of the reaction: samples of (*R*)-**9** rearranged to urea **13** in good yield without loss of er on both 200 mg and 1 g scales (Scheme 4).

Scheme 4. Stereospecific intramolecular arylation of an amine with a PMB (*p*-methoxybenzyl) protecting group.




Scheme 5. Stereospecific aryl migration within 2-pyridyl-stabilized anions.



2-Pyridyl-substituted organolithiums (2-picolylolithiums) have structures best characterized as azaenolates, as represented in Scheme 5 as **8Li-a**, with a planar α -carbon and the negative charge located principally at nitrogen.¹⁹ Urea-substituted enolates possessing other stereogenic centres undergo diastereoselective intramolecular arylation,²⁰ but (except for examples with special structural features promoting chiral memory by hindered rotation²¹) without stereospecificity at the planar enolate carbon atom.²² Nonetheless, treatment of the 2-pyridylurea **8a** with LDA in THF gave the rearranged compound **14a** with the same er (92:8) as the starting material (Scheme 5). Similarly, there was no loss of er in the rearrangements of ureas **8b** and **8c** bearing ethyl and allyl groups at the stereogenic centre: product ureas **14b** and **14c** were obtained in good yield and er (88%, 99:1 er and 67%, 94:6 er).

The stereospecificity of the α -(2-pyridyl)alkylamine synthesis was exploited by rearrangement of a range of *p*-methoxybenzyl-protected ureas **10a-h** built from chiral 2-pyridylamines (Scheme 6) to products **15a-h** without erosion of enantiomeric enrichment. Migrating rings with either electron-donating or withdrawing substituents at the *para* and *meta* positions all rearranged in high yield (85–98%) and good er (94:6–93:7) (**15b–15e**, **15g**). The migration of aromatic groups substituted at the *ortho* position gave lower yields: with 2-chlorophenylurea **10f** the reaction remained stereospecific but the yield dropped to 66%. Attempted rearrangement of a 1-naphthyl-substituent failed. Hydrolysis of the ureas **15** under basic conditions (NaOH, EtOH) gave the valuable 2-pyridyl substituted α -tertiary amines **16** in high yields (Scheme 6).²³

Scheme 6. Synthesis of protected tertiary α -(2-pyridyl)benzylamines by stereospecific intramolecular arylation. PMB = *p*-methoxybenzyl




We assume that in the rearrangements of ureas **7–10** the reaction mechanism proceeds by selective deprotonation of the acidic ‘picolinic’ position α to the pyridyl ring to give an anionic species **7–10Li**, which undergoes the conformational reorganisation required to attack the aryl ring borne by the other nitrogen atom, but without loss of configurational integrity. A non-classical intramolecular S_NAr reaction^{1b} leads to the product anion, which is protonated on work-up. The intermediate anionic species **7–10Li** must retain their absolute stereochemistry on the time scale of the rearrangement. This stereospecificity is a feature of the reaction shared with other related rearrangements of lithiated ureas^{9c,11a,24} (along with thiocarbamates²⁵ and, to a lesser extent, carbamates^{16,26}), though not the cyclic ureas so far explored.²⁷

The structure of 3-picolylolithiums related to **7–8Li** have not been examined in detail, but the inability of the nitrogen atom at the 3-position to stabilize the negative charge by delocalization suggests they may have structural similarities with configurationally stable α -nitrogen substituted benzyllithiums.³ By contrast, computational and crystallographic studies of 2-picolylolithiums related to **9–10Li**, whose negative charge is stabilized by delocalization onto the pyridyl nitrogen, show that the negative charge is principally located at the nitrogen atom, and the anion may be interpreted as a planar azaenolate, i.e. **8Li-a** (Scheme 5).¹⁹ Given the probable planarity of the α -carbon of the intermediate anion **8Li**, possible mechanisms for stereospecificity include long-lived planar chirality within an intermediate pyridyllithium species **8Li-b** (Scheme 5)^{25b–c,28} or the adoption by the urea of a chiral, twisted conformation (such as **8Li-c**) that rearranges to product faster than it can relax to an enantiomeric mixture of conformers.^{21,29} Attempts to probe the role of Li in the stereospecificity were frustrated by our inability to induce rearrangement with other bases (e.g. KHMDS).

Figure 3. In situ infra-red study of the rearrangement of **8b** to **11b**. 00:10:00 – Addition of 2.5 equiv LDA starts; 01:40:00 – Addition of 1 equiv LDA complete; 03:30:00 – Reaction quenched with MeOH.

(a)

(b)



Scheme 7. Proposed mechanistic pathway from **8b** to **14b**

To gain deeper insight into the mechanism of the reaction, the conversion of **8b** to **14b** in THF at $-78\text{ }^{\circ}\text{C}$ was followed by *in situ* infra-red spectroscopy (React-IR) (Figure 3 and Scheme 7). In THF at $-78\text{ }^{\circ}\text{C}$, IR shows one C=O stretching absorption at 1645 cm^{-1} and one pyridine C=N stretching absorption at 1594 cm^{-1} (Figure 3a). After 10 min (00:10) an initial 2.5 equiv LDA was added, causing the C=O stretch at 1645 cm^{-1} to diminish, the pyridine stretch to shift to 1591 cm^{-1} , and a new C=O stretch to grow at 1570 cm^{-1} . We assign these peaks to the rearranged, lithiated urea **C**.^{11a,30} Adding another equivalent of LDA 90 minutes later (01:40) completes the reaction, as indicated by the disappearance of the C=O stretch (starting material **A**) at 1645 cm^{-1} and a further increase of the C=O stretch (lithiated product **C**) at 1570 cm^{-1} . The detailed mechanism of formation of **C** from **B** was not explored, but previous studies^{11b} have suggested that related reactions proceed by a partially concerted $\text{S}_{\text{N}}\text{Ar}$ reaction. Finally (Figure 3b) addition of MeOH (03:20) protonates the urea anion of **C** to give **14b**, with a urea C=O stretch at 1633 cm^{-1} . No species identifiable as lithiated starting material **A** was observed. Product **14b** was recovered in 89% yield.

In summary, both α -(2-pyridyl) and α -(3-pyridyl) alkylamines may be arylated with total enantiospecificity at their α -position by intramolecular migration of an aryl substituent within their lithiated N' -aryl urea derivatives. Despite their delocalized structure, the intermediate 2-pyridyl-substituted anions are configurational stable on the time scale of the rearrangement.

ASSOCIATED CONTENT

Supporting Information

Full details of experimental procedures, characterization data and spectra of all new compounds; CIF file of (*R*)-**11c**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

†These authors contributed equally to this work. * Corresponding author e-mail: j.clayden@bristol.ac.uk.

ACKNOWLEDGMENT

This work was supported by the Marie Curie Intra-European Fellowship (IEF) “ENOLAR” and by the BBSRC. We thank Dr James Raftery (University of Manchester) for determining the X-ray crystal structure of (*R*)-**11c**.

REFERENCES

- (a) Roughley, S. D.; Jordan, A. M.; *J. Med. Chem.*, **2011**, *54*, 3451. (b) Carey, J. S.; Laffan, D.; Thomson, C.; Williams, M. T.; (c) Vitaku, E.; Smith, D. T.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 10257. (d) Yadav, A.; Verma, A.; Patel, S.; Kumar, A.; Rathore, V.; Meenakshi; Kumar, Sh.; Kumar, Sa. *Chem. Commun.* **2015**, *51*, 11658. (e) Goetz, A. E.; Garg, N. K. *Nature Chem.* **2013**, *5*, 54. (f) Neely, J. M.; Rovis, T. *Org. Chem. Front.* **2014**, *1*, 1010. (g) Rouquet, G.; Blakemore, D. C.; Ley, S. V.; *Chem. Commun.* **2014**, *50*, 8908.
- Taylor, R. D.; MacCoss, M.; Lawson, A. D. G., *J. Med. Chem.* **2014**, *57*, 5845.
- (a) Ullrich, T.; Krich, S.; Mereiter, K.; Anderson, D. J.; Meyer, M. D.; Pyerin, M. *J. Med. Chem.* **2002**, *45*, 4047. (b) Lawson, E. C.; Hoekstra, W. J.; Addo, M. F.; Andrade-Gordon, P.; Damiano, B. P.; Kauffman, J. A.; Mitchell, J. A.; Maryanoff, B. E. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 2619. (c) Kawata, S.; Ashzawa, S.; Hiram, M. *J. Am. Chem. Soc.* **1997**, *119*, 12012. (d) Rico, J. G.; Lindmark, J. R.; Bovy, P. R. *J. Org. Chem.* **1993**, *58*, 7948. (e) Aoki, M.; Ohtsuka, T.; Yamada, M.; Ohba, Y.; Yoshizaki, H.; Yasuno, H.; Sano, T.; Seto, H. *J. Antibiot.* **1991**, *44*, 582. (f) Atta-ur-Rahman; Shahwar, D.; Choudhary, M. I.; Sener, B.; Toker, G.; Baser, K. H. C. *Phytochemistry* **1999**, *50*, 333. (g) Husain, I.; Saquib, M.; Bajpai, V.; Kumar, B.; K. Shaw, A. K. *J. Org. Chem.* **2011**, *76*, 8930. (h) Bagley, M. C.; Dale, J. W.; Merritt, E. A.; Xiong, X. *Chem. Rev.* **2005**, *105*, 685. (i) Nicolaou, K. C.; Safina, B. S.; Zak, M.; Lee, S. H.; Nevalainen, M.; Bella, M.; Estrada, A. A.; Funke, C.; Zecri, F. J.; Bulat, S. *J. Am. Chem. Soc.* **2005**, *127*, 11159.
- (a) Haas, J.; Piguel, S.; Wirth, T. *Org. Lett.* **2002**, *4*, 297. (b) Brunner, H.; Markus, N. *Monatsh. Chem.* **2002**, *133*, 115. (c) Chelucci, G.; Pinna, G. A.; Saba, A. *Tetrahedron: Asymmetry* **1997**, *8*, 2571. (d) Canary, J. W.; Allen, C. S.; Castagnetto, J. M.; Wang, Y. *J. Am. Chem. Soc.* **1995**, *117*, 8484. (e) Chelucci, G.; Conti, S.; Falorni, M.; Giacomelli, G. *Tetrahedron* **1991**, *38*, 8251. (f) Brunner, H.; Heinrich, F. *J. Organomet. Chem.* **1987**, *335*, 1. (g) Zai, S.; Gao, H.; Huang, Z.; Hu, H.; Wu, H.; Wu, Q.; *ACS Catal.* **2012**, *2*, 433. (h) Baratta, W.; Benedetti, F.; Del Zotto, A.; Fanfoni, L.; Felluga, F.; Magnolia, S.; Putignano, E.; Rigo, P.; *Organometallics* **2010**, *29*, 3563. (i) Cheng, Y. Q.; Bian, Z.; Kang, C. Q.; Guo, H. Q.; Gao, L. X. *Tetrahedron: Asymmetry* **2008**, *19*, 1572. (j) Baratta, W.; Bosco, M.; Chelucci, G.; Del Zotto, A.; Siega, K.; Toniutti, M.; Zangrando, E.; Rigo, P.; *Organometallics* **2006**, *25*, 4611. (k) Yano, T.; Tanaka, R.; Nishioka, T.; Kinoshita, I.; Isobe, K.; Wright, L. J.; Collins, T. J. *Chem. Commun.* **2002**, 1396. (l) Rowland, J. M.; Olmstead, M. M.; Mascharak, P. K. *Inorg. Chem.* **2002**, *41*, 1545.
- Nguyen, T.; Shapiro, D. A.; George, S. R.; Setola, V.; Lee, D. K.; Cheng, R.; Rauser, L.; Lee, S. P.; Lynch, K. R.; Roth, B. L.; O'Dowd, B. F. *Mol. Pharmacol.* **2001**, *59*, 427.
- Eccles, R.; van Cauwenberge, P.; Tetzloff, W.; Borum, P. *J. Pharm. Pharmacol.* **1995**, *47*, 990.
- (a) Harikrishnan, L. S.; Finlay, H. J.; Qiao, J. X.; Kamau, M. G.; Jiang, J.; Wang, T. C.; Li, J.; Cooper, C. B.; Poss, M. A.; Adam, L. P.; Taylor, D. S.; Chen, A. Y. A.; Yin, X.; Sleph, P. G.; Yang, R. Z.; Sitkoff, D. F.; Galella, M. A.; Nirschl, D. S.; Van Kirk, K.; Miller, A. V.; Huang, C. S.; Chang, M.; Chen, X. Q.; Salvati, M. E.; Wexler, R. R.; Lawrence, R. M.; *J. Med. Chem.* **2012**, *55*, 6162. (b) Miller, M. M.; Liu, Y.; Jiang, J.; Johnson, J. A.; Kamau, M.; Nirschl, D. S.; Wang, Y.; Harikrishnan, L.; Taylor, D. S.; Chen, A. Y. A.; Yin, X.; Seethala, R.; Peterson, T. L.; Zvyaga, T.; Zhang, J.; Huang, C. S.; Wexler, R. R.; Poss, M. A.; Lawrence, R. M.; Adam, L. P.; Salvati, M. E.; *Biorg. Med. Chem. Lett.* **2012**, *22*, 6503.

- ⁸ Robak, M. T.; Herbage, M. A.; Ellman, J. A. *Chem. Rev.* **2010**, *110*, 3600.
- ⁹ (a) Shaw, A. W.; deSolms, S. J. *Tetrahedron Lett.* **2001**, *42*, 7173. (b) Spero, D. M.; Kapadia, S. R. *J. Org. Chem.* **1997**, *62*, 5537. (c) Clayden, J.; Hennecke, U. *Org. Lett.* **2008**, *10*, 3567.
- ¹⁰ Clayden, J. *Organolithiums: Selectivity for Synthesis*; Pergamon: Oxford, 2002.
- ¹¹ (a) Clayden, J.; Dufour, J.; Grainger, D. M.; Helliwell, M. *J. Am. Chem. Soc.* **2007**, *129*, 7488. (b) Vincent, M. A.; Maury, J.; Hillier, I. H.; Clayden, J. *Eur. J. Org. Chem.* **2015**, *5*, 953.
- ¹² Clayden, J. In *Lithium Compounds in Organic Synthesis*; Luisi, R., Capriati, V., Eds.; Dearomatization and Aryl Migration in Lithium Chemistry; 2014.
- ¹³ Kavirayani, R. P.; Omkar R. *Tetrahedron* **2013**, *69*, 8422.
- ¹⁴ Xing-Wen, S.; Min L.; Ming-Hua X.; Guo-Qiang L. *Org. Lett.* **2008**, *10*, 1259.
- ¹⁵ Chelucci, G.; Baldino, S.; Chessa, S.; Pinna, G. A.; Soccolini, F. *Tetrahedron: Asymmetry* **2006**, *17*, 3163.
- ¹⁶ Fournier, A. M.; Brown, R. A.; Farnaby, W.; Miyatake-Ondozabal, H.; Clayden, J. *Org. Lett.* **2010**, *12*, 2222.
- ¹⁷ Zimmerman, H. E.; Singer, L.; Thyagarajan, B. S. *J. Am. Chem. Soc.* **1959**, *81*, 108.
- ¹⁸ Clayden, J.; Donnard, M.; Lefranc, J.; Minassi, A.; Tetlow, D. J. *J. Am. Chem. Soc.* **2010**, *132*, 6624.
- ¹⁹ (a) Wiklund, T.; Olsson, S.; Lennartson, A. *Monatsh. Chem.* **2011**, *142*, 813. (b) Pratt, L. M.; Khan, I. M.; *THEOCHEM* **1996**, *367*, 33. (c) Holger, O.; Pieper, U.; Leusser, D.; Flierler, U.; Henn, J.; Stalke, D. *Angew. Chem. Int. Ed.* **2009**, *48*, 2978. (d) Pieper, U.; Stalke, D.; *Organometallics* **1993**, *12*, 1201.
- ²⁰ (a) Maury, J.; Clayden, J. *J. Org. Chem.* **2015**, *80*, 10757. (b) Atkinson, R. C.; Fernández-Nieto, F.; Mas Roselló, J.; Clayden, J. *Angew. Chem. Int. Ed.* **2015**, *54*, 8961.
- ²¹ Tomohara, K.; Yoshimura, T.; Hyakutake, R.; Yang, P.; Kawabata, T. *J. Am. Chem. Soc.* **2013**, *135*, 13294.
- ²² Atkinson, R. C.; Leonard, D. J.; Maury, J.; Castagnolo, D.; Volz, N.; Clayden, J. *Chem. Commun.* **2013**, *49*, 9734.
- ²³ The er of amines **16a-f**, like those of **12b,d** was not determined, but was assumed to be the same as the parent urea.
- ²⁴ Tait, M.; Donnard, M.; Minassi, A.; Lefranc, J.; Bechi, B.; Carbone, G.; O'Brien, P.; Clayden, J. *Org. Lett.* **2013**, *15*, 34.
- ²⁵ (a) MacLellan, P.; Clayden, J. *Chem. Commun.* **2011**, *47*, 3395. (b) Mingat, G.; MacLellan, P.; Laars, M.; Clayden, J. *Org. Lett.* **2014**, *16*, 1252. (c) Mingat, G.; McDouall, J. J. W.; Clayden, J. *Chem. Commun.* **2014**, *50*, 6754.
- ²⁶ Clayden, J.; Farnaby, W.; Grainger, D. M.; Hennecke, U.; Mancinelli, M.; Tetlow, D. J.; Hillier, I. H.; Vincent, M. A. *J. Am. Chem. Soc.* **2009**, *131*, 3410.
- ²⁷ (a) Bach, R.; Clayden, J.; Hennecke, U. *Synlett* **2009**, 421. (b) Tait, M. B.; Ottersbach, P. A.; Tetlow, D. J.; Clayden, J. *Org. Process Res. Dev.* **2014**, *18*, 1245. (c) Tait, M. B.; Butterworth, S.; Clayden, J. *Org. Lett.* **2015**, *17*, 1236.
- ²⁸ (a) Tetlow, D. J.; Hennecke, U.; Raftery, J.; Waring, M. J.; Clarke, D. S.; Clayden, J. *Org. Lett.* **2010**, *12*, 5442. (b) Tetlow, D. J.; Vincent, M. A.; Hillier, I. H.; Clayden, J. *Chem. Commun.* **2013**, *49*, 1548.
- ²⁹ Clayden, J.; Stimson, C. C.; Keenan, M.; Wheatley, A. E. H. *Chem. Commun.* **2004**, *2*, 228.
- ³⁰ (a) Grainger, D. M.; Smith, A. C.; Vincent, M. A.; Hillier, I. H.; Wheatley, A. E. H.; Clayden, J. *Eur. J. Org. Chem.* **2012**, *4*, 731. (b) Lefranc, J.; Fournier, A. M.; Mingat, G.; Herbert, S.; Marcelli, T.; Clayden, J. *J. Am. Chem. Soc.* **2012**, *134*, 7286.